




Male Contraception

Kirsten M. Vogelsong, PhD

UNDP/UNFPA/WHO/World Bank
Special Programme of Research
Development and Research Training
in Human Reproduction



Why Men in Family Planning?




- **International Conference on Population and Development, 1994**
- **Shared responsibility and gender equity**



Male Involvement in Fertility Regulation

- Condom
- Vasectomy
- Withdrawal
- Calendar/Rhythm





Distribution of Contraceptive Use Prevalence

World wide contraceptive use (Married Women of Reproductive age)

Contraceptive	No. of users (Millions)	Users (%)	First year failure rate (%) - Typical use
Total users	648	61.9	
<u>Modern methods</u>		56	
Female sterilization	210	20.1	0.5
IUD	156	14.9	0.8
Oral contraceptives	82	7.8	5.0
Condom	53	5.1	14.0
Male sterilization	43	4.1	0.15
Injectables	27	2.6	0.3
Vaginal barriers	4.2	0.4	20.0
<u>Traditional methods</u>			
Withdrawal	32	3.1	19.0
Rhythm	27	2.6	25.0

UN Population Division, 2001



Male Contraception

Research and Development

- Use of existing male methods is low, with regional and country differences
- Men are *aware* of family planning methods
- Men *approve* of the use of family planning
- Low levels of use may be related to the *negative characteristics of existing methods*
- Example: In a study conducted in Fiji, Iran, India and Korea, men considered a male pill or injection to be more acceptable than vasectomy




The Ideal Male Contraceptive

- **Safe** - *no harmful side effects*
- **Effective** - *it works!*
- **Acceptable** - *to men and their partners*
- **Affordable** - *to programs, potential users, and donors*



Approaches to Male Contraception: *Targeting the sperm*

- **Block deposition**
- **Interrupt transport**
- **Inhibit production**
- **Disrupt function**
- **Prevent fertilization**



Source: Image House Medical, Copenhagen



Blocking Sperm Deposition





Blocking Sperm Deposition

Male Condoms

- Condoms are effective at preventing pregnancy and STI/HIV
- Condom use is low even in countries with high prevalence of HIV/AIDS
- How can we increase condom use?



inSpiral



Blocking Sperm Deposition

Male Condoms

Condom studies

- Randomized comparative studies of “standard” and “new” condoms
 - Acceptability and preference
 - Contraceptive efficacy
 - Prevention of STI
- Reasons for use and non-use of condoms



Interrupting Sperm Transport

Vasectomy/Sterilization

World wide, nearly 43 million married couples rely on vasectomy

- United Kingdom - 18%
- New Zealand - 18%
- Canada - 15.2%
- Rep. of Korea - 13%
- United States - 13%
- The Netherlands - 11%
- Australia - 10%
- Switzerland - 8.3%
- Spain - 8.1%
- Bhutan - 8%
- China - 8%
- Belgium - 7.0%
- Nepal - 5.4%
- Thailand - 5.3%
- Denmark - 5%



Interrupting Sperm Transport

Vasectomy/Sterilization

- **Conventional vasectomy**
 - highly effective and safe
 - incision required
 - permanent
- **Percutaneous vas occlusion**
 - many compounds evaluated
 - lower efficacy rates
 - some additional complications
- **No-scalpel vasectomy**
 - highly effective
 - Somewhat more acceptable
 - lower complication rates



Methods of Vasectomy


Success of Reversal

Method	Follow-up (no and %)	Sperm (no and %)	Normal (no and %)	Pregnancy (no and %)
No-scalpel Vasectomy	19/23 (82.6)	16/19 (84.2)	13/19 (68.4)	15/19 (78.9)
Chemical Was occlusion	26/31 (83.9)	18/26 (69.2)	12/26 (46.2)	13/26 (50.0)
MPU Was occlusion	31/34 (91.2)	10/31 (32.3)	10/31 (32.3)	9/31 (29.0)



Inhibiting Sperm Production

Hormonal Contraception





Inhibiting Sperm Production

Hormonal Contraception

Androgen alone

T Enanthate

T Undecanoate

T Buciclate

Pellets

Progestin + Androgen

Norplant

DMPA

Norethisterone Enanthate

GnRH

Agonists

Antagonists

Vaccines

FSH

Antagonists

Vaccines



Hormonal Approaches to Male Contraception


Androgen alone

- 1990: 200 mg testosterone enanthate/week will reduce sperm production in some men
- Sperm concentrations consistently below 1 million/ml result in few or zero pregnancies
- All men do not fully suppress
- Requirement for weekly injections and high T concentrations



Hormonal Approaches to Male Contraception

Androgen alone




Sperm concentrations following weekly inj. 200 mg T-enanthatate



Hormonal Approaches to Male Contraception

Androgen alone



Testosterone Enanthate

- Extensive clinical experience
- “Burst” effect
- Short acting
- Weekly injections
- High levels testosterone



Hormonal Approaches to Male Contraception

Androgen alone

Testosterone Undecanoate

- Oral or injectable
- Longer release profile
- 4-8 week injection intervals may be adequate
- Maintains testosterone in physiological range
- Large dose required


Testosterone Buciclate

- No “burst” effect
- Long-acting injectable
- Injections at 3-4 month intervals may be adequate
- High dose required



Hormonal Approaches to Male Contraception

Androgen alone





Hormonal Approaches to Male Contraception

Androgen with Progestin

- More rapid and effective sperm suppression
- Effective in diverse populations
- Reduced overall drug load
- Physiological testosterone levels
- Requires a 2 drug regimen
- Drugs may have different routes or frequencies of administration



Hormonal Approaches to Male Contraception

Androgen with Progestin

Progestagen	Androgen	% Azoo- spermic	% Oligozoo spermic	Reference
DMPA 250 mg every 6 weeks	19 NT (200 mg every week x 6/7 weeks, then 200 mg/3 or 4 weeks).	67 (W) 98 (A)	92 (W) 99 (A)	Knuth et al (1987)
	TE (200 mg(IM every week x 6/7 weeks, then 200 mg/4 weeks)	59 (W) 96 (A)	91 (W) 96 (A)	WHO (1993)
DMPA 300 mg	T implant (800 mg)	90 (W)	100 (W)	Handelsman et al (1996)



Hormonal Approaches to Male Contraception

Androgen with Progestin


Progestagen	Androgen	% Azoo-spermic	% Oligozoo-spermic	Reference
Levonorgestrel (oral)				
500 µg/day	TE (100 mg/week IM)	67 (W)	94 (W)	Bebb et al (1996)
250 µg/day	TE (100 mg/week IM)	78 (W)	89 (W)	Anawalt et al (1997)
125 µg/day	TE (100 mg/week IM)	61 (W)	94 (W)	
Desogestrel (oral)				
300 µg/day	TE (100 mg/week IM)	81 (W)	94 (W)	Wu et al (1998)
150 µg/day	TE (50 mg/week IM)	73 (W)	100 (W)	

W=White, A=Asian, DMPA=depotmedroxyprogesterone acetate, TE=testosterone enanthate
19 NT= 19 nortestosterone hexyloxyphenylpropionate



Hormonal Approaches to Male Contraception

Androgen with Progestin





Hormonal Approaches to Male Contraception

Other Approaches

- Androgen with anti-androgen (*cyproterone acetate*)
 - Progestin with anti-androgen properties
 - May block the activity of any residual T in the testis
- Androgen with GnRH Analogue
 - Effective suppression of gonadotrophins
 - High cost; frequent application



Disrupting Sperm Function and Preventing Fertilization

- Targeted basic science research on testicular, epididymal or vas approaches

Some promising targets:

- functional development, i.e. motility
- structural development, i.e. organelles
- structure and function, i.e. membrane integrity and intracellular pathways



Male Reproductive Health Agenda

- **Contraceptive research and development**
- **Targeted basic science -physiology and fertility**
- **Social & behavioral sciences**
- **Men's roles in reproductive health**
- **Building networks**



Acceptability/Sociobehavioral Studies

- Current use of male methods
- Preferences for new methods
- Characteristics of new methods
- Continuation and discontinuation of trial
- Effects on mood
- Effects on behavior
- Effects on cognition
- Partner's views on mood and behavior



Acceptability/Sociobehavioral Studies

Reports from 25 Swedish men participating in TE trial

Expectations

- Freedom and security
- Problems with female methods
- Desire for more satisfying sex life
- Need for male control
- Fear of negative side effects

Satisfaction

- Greater freedom
- More ease in sex life
- Would recommend method to others
- Trouble with injections
- Fear of problems with aggressiveness
- Dermatological problems



Acceptability/Sociobehavioral Studies

	<i>Very important</i>	<i>Somewhat important</i>	<i>Not important</i>
Men should share responsibility for contraception	41.2	51.0	7.8
Contributing to solving the population problem	41.6	48.7	9.7
I felt I was doing a good thing for my country	36.7	52.9	7.9
I like to be involved in new things	25.0	56.8	18.2
I felt pride in contributing to scientific advancement	26.9	51.6	21.4
Pioneer of a new method of contraception	24.4	46.1	29.5
My wife wanted me to take responsibility	23.1	44.8	32.1
I joined for getting the financial compensation	12.7	28.6	58.8



Acceptability/Sociobehavioral Studies

	Month 4	Month 8
	%	%
Reasons for perceived inconvenience	(n = 78)	(n = 117)
Have to come to clinic	23.1	9.3
Once a month too frequent	70.5	76.3
Wait at the clinic	1.3	5.1
Other	5.1	9.3
Total	100.0	100.0
Reasons for dissatisfaction	(n = 87)	(n = 117)
Side effect	11.5	6.0
Inconvenience	54.0	48.7
Injection pain	21.8	12.0
Others	12.6	33.3
Total	100.0	100.0



Men's Roles in Reproductive Health

Men can:

- **Inhibit access to and use of FP**
 - **Expose women and themselves to disease including HIV**
 - **Act as barriers to women's reproductive health**
- OR**
- **Facilitate & support use of contraception**
 - **Protect themselves and their partners from infection**
 - **Act as partners in promoting reproductive rights and care for all**



Providing FP Services to Men

- How can FP service facilities address men's needs?
- How to create and then address an increase in demand for FP services for men?
- Who will provide FP services to men?