Spontaneous Abortion Recurrent Pregnancy loss

Dr. Anan Sacdpraseuth Mahosot Hospital

GFMER - WHO - UNFPA - LAO PDR
Training Course in Reproductive Health Research
Vientiane, 26 November 2009

Five Key Points

- 1. Recurrent pregnancy loss occurs in 2% to 5% of all couples of reproductive health.
- 2. Increased maternal age is a significant risk factor.
- 3. Causes include genetic, endocrinologic, anatomic, immunologic, infectious, psychological, lifestyle and environmental factors.
- 4. When the physician decides that an evaluation is warranted, a complete diagnosis evaluation should be initiated.
- 5. Alcohol consumption (> 5 drinks per week) and tobacco use (>10 cigarettes per day) increase the occurrence of pregnancy loss 1.5 to 2 fold over the age-adjusted risk.

Spontaneous Abortion Definition

- Non induced abortion of product of conception before 20th weeks of pregnancy.
- Natural termination of pregnancy before the fetus is capable of external life.

Diagnosis of Abortion DIAGNOSIS BLEEDING CERVIX UTERINE SIZE

Dilated

Dilated or

closed

Closed

OTHER SIGNS

Positive pregnancy

Uterus tender/firm

Partial expulsion of

Uterus tender/firm

Complete expulsion

of products of

Fetus dead with

delayed expulsion

pregnancy signs and

conception

Decrease in

symptoms

test

Less than or

Less than

Less than or

equal to dates

dates

equal to dates

Cramping

Uterus soft

Cramping

products of conception

l to dates	Equal to	Not dilated	Slight to moderate	Threatened Abortion
				er .
than or to dates	Less tha	Dilated	Moderate heavy	Inevitable Abortion

Slight to heavy

Slight to

moderate

Little or none

Incomplete

Abortion

Complete

Abortion

Missed

Abortion

Occurrence and Recurrence

- 10 to 15 % of recognized pregnancies.
- About 20 to 30% of women with confirmed pregnancies bleed during the 1st 20 wk of pregnancy; 1/2 of these women spontaneously abort.
- Clinical abortion in women aged over 40 is more than double the rate in women younger than 20.

Maternal Age	Percent Abortion	Paternal Age	Percent Abortion
<20	12.2	<20	12
20-24	14.3	20-24	11.8
25-29	13.7	25-29	15.7
30-34	15.5	30-34	13.1
35-39	18.7	35-39	15.8
40-44	33.8	40-44	19.5
>44	53.2	>44	na

Prior outcome	Prior losses	Recurrence%
Liveborn	0	12
	1	24
	2	26
	3	32
	4	26
Miscarriage	1	19
	2	35
	3	47
	4	54