

# Updates on Contraceptive Technology



Training Course in Sexual and Reproductive Health Research  
Geneva 2011

08\_XXX\_MM1

# Updates on Contraceptive Technology Part 2

Kirsten Vogelsong  
Mario Festin  
Promoting Family Planning  
Department of Reproductive Health and Research





Training Course in Sexual and Reproductive Health Research  
Geneva 2011

08\_XXX\_MM2

# Comparing Effectiveness of Family Planning Methods

More effective

Less than 1 pregnancy per 100 women in one year



## How to make your method more effective

**Implants, IUD, female sterilization:**

After procedure, little or nothing to do or remember

**Vasectomy:** Use another method for first 3 months

**Injectables:** Get repeat injections on time

**Lactational Amenorrhea Method (for 6 months):** Breastfeed often, day and night

**Pills:** Take a pill at the same time each day

**Patch, ring:** Keep in place, change on time

**Condoms, diaphragm:** Use correctly every time you have sex

**Fertility awareness methods:** Abstain or use condoms on fertile days. Standard Days Method and Two-Day Method may be easier to use.

**Withdrawal, spermicides:** Use correctly every time you have sex

Less effective

About 30 pregnancies per 100 women in one year

08\_XY

# Outline and Objectives

- Description of the method
- Mechanism of action
- Effectiveness
- Eligibility criteria
- Benefits and side effects
- Interventions for associated effects

08\_XXX\_MM4

# Methods

- Combined oral contraceptives
- Progestin only pills
- Injectable contraceptives, progestin-only
- Injectable contraceptives, combined
- Hormonal implants
- **IUDs (copper bearing)**
- **LNG IUS**
- **Male and female condoms**
- **Other barrier methods**
- **Fertility awareness, lactational amenorrhea**
- **Emergency contraception**
- **Tubal ligation and vasectomy**
- **Other methods**


# Copper Intrauterine Contraceptive Devices (IUDs or IUCDs)



Copper T 380A


08\_XXX\_MM6

# What is a Copper IUD?



- Small plastic device with copper sleeves or wire around it, inserted through the vagina and cervix into the uterus
  - Safe
  - Highly effective
  - Long acting (up to 12 years)
  - Require trained provider to insert and remove

# Copper-bearing IUD



Copper T-380A

TCu-380A, “Copper T” is most widely used copper IUD



Multiload 375


Multiload 375



# Copper T: Mechanism of Action

Prevents fertilization by:

- Impairing the viability of the sperm
- Interfering with movement of the sperm



Source: Ortiz, 1996

# Copper IUDs: Characteristics

- Safe and highly effective
- Require no user action
- Long-acting (up to 12 years)
- Rapid return to fertility
- No systemic effects
- Other health benefits
- Potential side effects
- Require pelvic exam
- Trained provider needed to insert and remove
- Possible pain or discomfort during insertion
- Do not protect against STIs/HIV
- Rare complications

Source: CCP and WHO, 2011

# Copper IUDs: Possible Side Effects

Some women may experience:

- More cramps and pain during menstruation
- Increased or prolonged menstrual bleeding
- Bleeding between menstrual periods



***Side effects are most common during the first 3–6 months.***

Source: CCP and WHO, 2011; Larsson, 1993; DeMaeyer, 1989; WHO, 2004, updated 2008; WHO Special Programme of Research Development and Research Training in Human Reproduction, 1997

# Copper IUDs: Health Benefits

IUDs are known to:

- Reduce risk of ectopic pregnancy
  - Rate in all IUD users is 12 in 10,000 (2 in 10,000 for Copper T380A users)
  - Rate in women using no contraception is 65 in 10,000
- Help protect against endometrial cancer

Source: CCP and WHO, 2011; Sivin, 1991.

# Potential Complications

- Perforations—Very rare, 1 in 1,000 insertions
  - Linked to skill and experience of provider
  - Reduced through supervised training
- PID—Rare, most due to gonorrhoea or chlamydia at time of insertion
- Expulsions—Related to provider skill, age and parity of woman and insertion factors

08\_XXX\_MM13

# Expulsion Rates Are Higher for Postpartum Insertion

Timing of Insertion	Expulsion Rates
<b>Interval</b> (more than 6 weeks after delivery)	<b>Low</b> (3% for skilled provider)
<b>Immediate postpartum</b> (within 10 minutes)	<b>Slightly higher</b>
<b>Early postpartum</b> (between 10 minutes and 48 hours)	<b>Moderately higher</b>
<b>Late Postpartum</b> (48 hours to 4 weeks)	<b>High - Generally not recommended</b>

Source: Chi, et al, 1985.

# Who Can Initiate Copper IUDs

WHO MEC category 1 and 2 examples

WHO MEC Category	Conditions (selected examples)
Category 1	$\geq 20$ years, cervical ectopy, uterine fibroids without distortion of the uterine cavity, irregular bleeding without heavy bleeding, breastfeeding $> 6$ weeks, history of PID
Category 2	Menarche to $< 20$ years, nulliparous, heavy or prolonged bleeding, severe dysmenorrhea, anemia, HIV/AIDS infection, with or without ARV therapy

Source: WHO, 2004; updated 2008.

# Who Should Not Initiate Copper IUDs

WHO MEC category 3 and 4 examples

WHO MEC Category	Conditions (selected examples)
Category 3	48 hours to <4 weeks postpartum, ovarian cancer/if initiating use, high individual risk of STI/ if initiating use
Category 4	pregnancy, unexplained vaginal bleeding (prior to evaluation), current PID or cervical infection, endometrial or cervical cancer/if initiating use

Source: WHO, 2004; updated 2008.



# Timing of IUD Insertion

## Interval insertion

- Within the first 12 days of menstrual cycle
- Any other time if woman is not pregnant

## Postpartum insertion

- Immediately after vaginal or cesarean delivery if no infection or bleeding complications
- Within 48 hours or delay at least 4 weeks

## Postabortion insertion

- Immediately or within 12 days if no infection

Source: WHO, 2004; updated 2008.

# Side Effects

- Common side effects:
  - Heavier and/or prolonged menstrual bleeding
  - Menstrual cramping
  - Spotting between periods
- Side effects:
  - Are not signs of illness
  - Usually become less within the first 3–6 months

# Correcting Misconceptions

## IUDs:

- Rarely lead to PID
- Do not increase risk of STIs, including HIV
- Do not work by causing abortion
- Do not make women infertile
- Do not move to the heart or brain
- Do not cause birth defects
- Do not cause pain for either partner during sex
- Significantly reduce risk of ectopic pregnancy

08\_XXX\_MM19



# Signs of Possible IUD Complications

- Bleeding and severe abdominal cramping within a few days post-insertion ➡ perforation
- Irregular bleeding or pain every cycle ➡ partial expulsion, perforation
- Fever, unusual vaginal discharge, low abdominal pain ➡ infection
- Missing IUD strings, missed period ➡ expulsion, pregnancy

Source: CCP and WHO, 2011.

Managing IUD Side Effects or Complications:  
**Heavy, Prolonged or Irregular Bleeding**

***Counseling and reassurance are key***

<b>Problem</b>	<b>Action/Management</b>
<b>Heavy or prolonged bleeding</b>	<ul style="list-style-type: none"><li>● Reassure client that this is common and not harmful, usually diminishes after few months</li><li>● For short-term relief offer 5-day course of tranexamic acid or NSAIDs (not aspirin)</li><li>● Provide iron tablets</li></ul>
<b>Irregular bleeding</b>	<ul style="list-style-type: none"><li>● Reassure client</li><li>● For short-term relief offer ibuprofen or indomethacin 2 times daily after meals for 5 days</li></ul>

08\_XXX\_MM21

Source: CCP and WHO, 2011.

# Managing IUD Side Effects or Complications: Cramping and Mild Pain

***Counseling and reassurance are key***

Problem	Action/Management
<b>Cramping and mild pain</b>	<ul style="list-style-type: none"><li>● She can expect cramping and pain in first 1–2 days after insertion</li><li>● Reassure client that this is common in first 3–6 months, not harmful, usually decreases over time</li><li>● Suggest ibuprofen, other pain reliever (not aspirin if she also has heavy bleeding)</li><li>● If cramping continues, occurs outside of menstruation, evaluate, treat or refer</li><li>● If cramping is severe but no underlying condition, discuss removing the IUD</li></ul>

Source: CCP and WHO, 2011.

## Managing IUD Side Effects or Complications: Severe Pain in Lower Abdomen (Rare)

- Rule out PID, ectopic pregnancy or perforation.
- If PID is suspected, treat with appropriate antibiotics for gonorrhoea, chlamydia and anaerobic bacterial infection. There is no need to remove the IUD.
- If ectopic pregnancy is suspected, refer immediately.

Source: CCP and WHO, 2011.

## Managing IUD Side Effects or Complications: Suspected Perforation

- Stop procedure immediately, remove IUD
- Observe vital signs for an hour; check for signs of bleeding
  - If rapid pulse, falling blood pressure, or increased pain: refer
- Provide alternative contraception
  - Advise avoid sex for 2 weeks
- Follow-up in a week or as needed

Source: CCP and WHO, 2011.



## Managing IUD Side Effects or Complications: Missing Strings

- Determine risk of pregnancy
- Perform pelvic exam, probe for strings in cervical canal
- If cannot locate strings, consider X-ray or ultrasound, or refer
- Give choice of another contraceptive method
- Insert another IUD if expulsion is confirmed and
  - Woman is not pregnant
  - She still wants to use an IUD

Source: CCP and WHO, 2011.

## Managing IUD Side Effects or Complications:

# Suspected Pregnancy

- Assess for pregnancy, including ectopic pregnancy
- If the client is pregnant and wishes to continue the pregnancy:
  - Explain that using an IUD during pregnancy increases the risk of preterm delivery or miscarriage
  - If possible, remove the IUD
  - If not possible to remove, advise close follow-up for signs of septic miscarriage

Source: CCP and WHO, 2011.

# Hormonal intrauterine device



LNG-IUS

08\_XXX\_MM27

# Hormonal IUDs




- Steadily release a progestin
- Levonorgestrel IUD (LNG-IUD) most common hormonal IUD
- Also known as LNG-intrauterine system (LNG-IUS)
- Immediately reversible
- Effective for 5 years

# Other features of the LNG IUS

- Works primarily by suppressing the growth of the lining of the uterus
- Needs to be inserted into a uterus by a trained health care provider
- Some report lighter, fewer or even absent bleeding days
- Also infrequent reports of headaches, breast tenderness or pain, acne

08\_XXX\_MM29


# Male and Female Condoms



08\_XXX\_MM30

# What Is a Male Condom?

- A sheath that fits over a man's erect penis
- Most are made of thin latex rubber
- Other materials include natural skin and various synthetics



08\_XXX\_MM31



# What Is a Female Condom?

- A transparent sheath that fits loosely inside a woman's vagina
- Most common type is FC2—made of thin, soft, synthetic rubber film
  - Has flexible rings at both ends
- Natural latex female condoms available in some countries



08\_XXX\_MM32



# Comparing Effectiveness of Male and Female Condoms

*In real-life situations, correct and consistent use may be difficult to achieve*

## Pregnancy rates:

Male



Female




perfect use	2%	5%
typical use	15%	21%


Source: Hatcher, 2007.

# Condoms: Mechanism of Action

- **Create a barrier that keeps sperm from entering the vagina, thus preventing pregnancy**
- **Also form a barrier against STIs including HIV**



Male and Female Condoms  
**Dual Protection**



Condoms are the *only* FP method that provides **dual protection**:

- Protection from pregnancy *and*
- Protection from transmission of HIV and other STIs between partners

# Effectiveness for Preventing HIV and STIs

- The consistent, correct use of condoms significantly reduces the risk of HIV infection in men and women
- When used correctly with every act of sex, condoms are 80%–95% effective in preventing HIV infection
- Condoms prevent STIs transmitted through body fluids such as gonorrhea and chlamydia
- Condoms are less effective for preventing STIs transmitted by skin-to-skin contact, such as herpes and warts

# Characteristics of Male Condoms



- Safe and easy to use
- Widely available
- Effective when used consistently and correctly
- Provide dual protection
- No hormonal side effects
- Can help men with premature ejaculation
- Do not require provider's help
- Can be used as temporary backup method
- Protect women from conditions caused by STIs

08\_XXX\_MM37

# Characteristics of Male Condoms

- As typically used, less effective than many other methods
- Require partner communication and cooperation
- Can be damaged by oil-based lubricants, heat, humidity or light
- May reduce sensation
- Can interrupt sex





# Who Can Use Male Condoms

- All men and women can safely use male condoms, *except* those with severe allergic reaction to latex
  - Extremely rare among both men and women
  - Non-latex condoms are available in some countries



## Male Condoms

# Managing Problems

- If condom not used—Offer emergency contraception
- If slipping or breaking—Ask about practices, behaviors
- If difficulty putting on—Ask client to demonstrate, correct errors
- Difficulty persuading partner to use—Help her choose approaches that will work  
Suggest adding another method





## Male Condoms

# Managing Problems

Mild irritation or mild allergic reaction to condom—Itching, redness, rash and/or swelling

- Try another brand of condoms
- Put lubricant or water on the condom
- If symptoms continue, assess or refer for possible vaginal infection or STI
- If no infection, may have allergy to latex

# Characteristics of **Female Condoms**

## What Men and Women Like

### What Women Like:

- Female-controlled
- Texture feels more natural than latex male condoms
- Offer STI/HIV protection
- Outer ring provides stimulation
- Do not require provider's help

### What Men Like:

- Can be inserted in advance
- Are not tight or constricting
- Do not dull sensation
- Do not have to be removed immediately

08\_XXX\_MM42

## Characteristics of **Female Condoms**

# What Men and Women Don't Like

### **Limitations:**

- May be difficult to insert at first, require practice
- Not as effective as other methods
- More expensive than male condoms
- Less available than male condoms

08\_XXX\_MM43

# Who Can Use Female Condoms


- All men and women can use FC2
- Latex female condoms: Severe allergic reaction to latex is the only condition that prevents use
  - Extremely rare



08\_XXX\_MM44

# Managing Problems

- If having trouble inserting: Ask her to demonstrate
- If uncomfortable: Reinsert or reposition
- If condom squeaks: Use more lubricant
- If condoms slips or is not used correctly: Offer emergency contraception
- Difficulty persuading partner: Help her choose approaches that will work



## Female Condoms

# Managing Problems

(Continued)

- Mild irritation in or around the vagina or penis (itching, redness, rash)
  - Usually goes away on its own
  - Suggest added lubricant inside condom or on penis
  - If symptoms persist, assess and treat for possible vaginal infection or STI

# Responding to Myths and Fears

## Male condoms:

- Do not make men impotent, weak, or decrease sex drive
- Do not cause illness in men because sperm “backs-up”

## Female condoms:

- Are not difficult to use, but correct use needs to be learned
- Do not cause illness in women because they prevent semen from entering the body



# Responding to Myths and Fears

(continued)

## **Both male and female condoms:**

- Cannot get lost in woman's body
- Do not have holes that HIV can pass through
- Are not intended only for use outside of marriage, but also used by married couples

08\_XXX\_MM48



# Other barrier methods

- **Diaphragm** – is placed deep in the vagina before sex, covering the cervix as a barrier, usually used with a spermicide
- When used correctly with every act of sex, 6 pregnancies per 100 women using the diaphragm over the first year.
- **Cervical caps** – is also placed deep in the vagina before sex, covering the cervix as a barrier.
- Not as effective, with 20 pregnancies per 100 women in the first year.



# Other barrier methods

- Spermicide – applied deep in the vagina before every act of sex
  - Usually Nonoxynol 9, but other substances also available
  - Work by causing the membrane of sperm cells to break, killing them or slowing movement.
  - One of least effective methods, may be a primary or a back up method.
- Microbicide (either anti-infective alone or dual protection)



08\_XXX\_MM50

# Fertility awareness methods

- Requires the cooperation of both partners, with a commitment to abstain or use another method on fertile days
  - Periodic abstinence
  - Natural family planning
- Must be aware of the body changes or keep track of days, according to rules of the specific methods.
- No side effects.

# Fertility awareness methods

- Calendar based methods
  - Standard days methods
  - Calendar rhythm methods
- Symptoms based methods
  - Cervical secretions
  - Basal Body Temperature
    - Increase noted slightly after release of the egg or ovulation
  - Two day method
  - Sympto-thermal method

08\_XXX\_MM52


# Standard Days Methods

- 95% effective with consistent and correct use
- A woman can use the SDM if most of her menstrual periods are 26 to 32 days long.
- Days 8 to 19 of each cycle are considered as fertile days.
- The couple can use cycle beads, a color-coded string of beads that indicates fertile and nonfertile days.



08\_XXX\_MM53

# How to use cycle beads



- Move ring to RED bead when period starts

1



- Move ring to next bead every day. Move ring even on bleeding days

2



- Use condoms or abstain when ring is on WHITE beads

3




- BROWN beads are safe days of no pregnancy

4



- When period starts again move ring to red bead to begin again.

5



- Always check your period comes between dark brown bead and last brown bead.



# Lactational amenorrhea

- A family planning method based on breastfeeding
- Can be effective for up to 6 months after delivery, as long as monthly bleeding has not returned and the woman is fully breastfeeding.
- Provides an opportunity to offer a woman an ongoing method for continuously 6 months

08\_XXX\_MM55



# Lactational amenorrhea

- A woman with the following may want to consider other methods:
  - Has HIV infection
    - Especially if the woman is not taking ARVs, not fully breastfeeding, newly infected or has advanced disease.
  - Using certain medications (mood-altering drugs, reserpine, ergotamine, anti-metabolites, cyclosporine, high dose corticosteroids, bromocriptine, radioactive drugs, lithium and certain anticoagulants)
  - A newborn with a condition that makes it difficult to breastfeed



# Emergency contraception

- Emergency contraceptive pills help to prevent pregnancy, if taken up to 5 days after unprotected sex and other situations such as
  - forced sex (rape) and contraceptive mistakes (condom slippage or breakage, misplaced IUD, late for injections, etc)
- Do not disrupt an existing pregnancy
- Safe for all women
- Provide an opportunity for women to start using an ongoing FP method
- Effectiveness (no method = 8 pregnancies/100 woman years)
  - Progestin only ECPs – 1 pregnancy per 100 women years
  - Combined estrogen progestin ECPs – 2 pregnancies per 100 women years.
- Copper IUD effective as emergency contraception, when inserted up to 7 days after unprotected sex

08\_XXX\_MM57

# Emergency contraception

- Contains a progestin alone (levonorgestrel or ulipristal) or a progestin and an estrogen together
- Works primarily by preventing or delaying ovulation
- Does not work if the woman is already pregnant.

08\_XXX\_MM58

<b>Pill type</b>	<b>Total dosage to provide</b>
<b>Levonorgestrel-only dedicated product</b>	<ul style="list-style-type: none"> <li>• 1.5 mg of levonorgestrel in a single dose.<sup>6</sup></li> </ul>
<b>Estrogen-progestin dedicated product</b>	<ul style="list-style-type: none"> <li>• 0.1 mg ethinyl estradiol + 0.5 mg levonorgestrel. Follow with same dose 12 hours later.</li> </ul>
<b>Progestin-only pills with levonorgestrel or norgestrel</b>	<ul style="list-style-type: none"> <li>• Levonorgestrel pills: 1.5 mg levonorgestrel in a single dose.</li> <li>• Norgestrel pills: 3 mg norgestrel in a single dose.</li> </ul>
<b>Combined (estrogen-progestin) oral contraceptives containing levonorgestrel, norgestrel, or norethindrone</b>	<ul style="list-style-type: none"> <li>• Estrogen and levonorgestrel pills: 0.1 mg ethinyl estradiol + 0.5 mg levonorgestrel. Follow with same dose 12 hours later.</li> <li>• Estrogen and norgestrel pills: 0.1 mg ethinyl estradiol + 1 mg norgestrel. Follow with same dose 12 hours later.</li> <li>• Estrogen and norethindrone pills: 0.1 mg ethinyl estradiol + 2 mg norethindrone. Follow with same dose 12 hours later.</li> </ul>

# Correcting misconceptions about Emergency contraception


ECPs :

- Do not cause abortion
- Do not cause birth defects if pregnancy occurs
- Are not dangerous to a woman's health
- Do not promote sexual risk taking
- Do not make women infertile

08\_XXX\_MM60

# Female Sterilization Tubal Ligation


- Permanent method, done through mini-laparotomy (small incision) or laparoscopy
- One of the most effective methods with less than 5 failures per 1000 women
- May be used by any woman
- Counselling needed to ensure no post procedural regrets



08\_XXX\_MM61

# Male Sterilization

## Vasectomy



- Simple permanent surgical method in men
- With a 3 month delay in taking effect due to storage of sperm
- Tubes that carry sperm to the penis (vas deferens) are cut and ligated (by sutures or cautery)
- May be done using non-scalpel technique
- Less than 2 per 1000 pregnancies over first year
- Does not affect male performance nor provides increase risk for cancer




# Other methods

- Ring

- Requires keeping a flexible ring which releases progestins and estrogen which are absorbed by the body
- Kept for 3 weeks, and woman menstruates in 4<sup>th</sup> week
- Prevents ovulation

- Patch

- Requires wearing a small adhesive patch with estrogen and progestins weekly, for 3 weeks
- Works by preventing ovulation



# Acknowledgments and References

- Main Reference:
  - Family Planning – A Global Handbook for Providers (<http://www.globalhandbook.org/>)
- Acknowledgements
  - Family Health International
  - Knowledge for Health
  - Institute of Reproductive Health



# Updates on Contraceptive Technology



08\_XXX\_MM65