

Training Course in Sexual and Reproductive Health Research 2013 Module: Principles and Practice of Sexually Transmitted Infections Prevention and Care

Improving M&E STI/HIV programmes


Keith Sabin - UNAIDS

with

WHO	UNDOC	UNC/MEASURE
Martin Donoghoe	Riku Lehtovuori	Sharon Weir
David Jacka	Mirlan Mamyrov	









Roles of M&E of STI Program

- Global
 - Assess progress toward universal access to STI/HIV prevention, care and treatment
 - Assess global response across countries
- National
 - Assess national program response to STI/HIV
- Local
 - Is our project working? Are we reducing STI/HIV?



3 Stepped Approached


- Agree on useful indicators
- Set targets for accepted indicators
- Collect data on indicators



3 Steps Elaborated

- Agree on useful indicators
 - What indicators will improve program?
- Set targets for accepted indicators
 - Use Technical Guide
- Collect data on indicators
 - Use operational guideline





Data Management for M&E

Oh yeah, and report the data!

- National programs should have information management systems sufficient to:
 - Aggregate,
 - Analyze,
 - Interpret,
 - and report the data.

If you don't use the data, it's not worth collecting it!!!!



Baby Steps

Define, define, define

- What is the case definition of each STI?
 - Lab-based or syndromic?
- What is the definition of key populations?




Baby Steps

- Define, define, define
- What is the case definition of STI?
 - What client information will you collect?
 - What ages?
 - What geographic areas?
 - How recently?
 - Risk behaviors?
 - Quality of experience using service?



Target Setting Guide*

- "Represents an initial attempt to develop a framework for assessing countries' progress in delivering a comprehensive package of core interventions."
- For each service in the comprehensive package, Guide suggests indicators for:
 - Availability
 - Coverage
 - Quality
 - Impact
- But.. "Guidance on operationalization, and monitoring and evaluation of interventions is outside the scope of this Guide..." - Operational guideline to respond to this need
- Revised in 2012.





Indicators

- Selecting useful indicators
 - Steps to choosing data elements to collect

You are the manager of a needle and syringe program.

The National STI/HIV Program asks you to report number of condoms distributed per sex worker per year.

- 1. How do you do this? Do you want to know this?
- 2. What would you want to know?
- 3. How do you decide?



Targets

Do we need targets for our program? Why?
 How do we use them? How do we set them?



What is the current programmatic response and what are the targets?

The objectives of this step

- To define the package of services that should be provided as part of the national and sub-national response
- To map the availability of services that are part of the comprehensive package in each subnational area
- To specify coverage, outcome, and impact indicator targets for the national level and in each sub-national area



3.1 Based on the epidemic among the population, what services are needed in response?



3.2 In each sub-national area:

-What services are currently available?

-What contributing factors are being addressed?



3.3 Based on the assessment, what are the 2-year targets for impact, outcome and coverage indicators in each sub-national area?



How to Answer the Key Questions


Key Question	Methods	s Products	Use
3.1 Based on the epidemic ampeople who sex and analy of contributing factors, what combination services and activities is needed to prevent STI transmission	Planning rsis ng of	Documents that define the combination of services and activities provided as part of the national response	To define coverage indicators (see Step 6)



How to Answer the Key Questions

	Key Questions	Methods	Products	Use
3.2	What services are currently provided in each sub-national area? What other activities are being undertaken to address factors that contribute to the HIV epidemic?	1. Service availability mapping and Stakeholder Meetings	Map showing geographic availability of services and community-strengthening activities	To monitor service availability and identify gaps in services

STI intervention sites for high-risk groups





Target Setting Steps

	Key Questions	Methods	Products	Use
3.3	What are the current baseline and the two year targets for changes in coverage, outcome and impact indicators?	1. Target setting methods	Completed target- setting worksheet	Assess programme effectiveness (see Step 7)



Targets are:

- Set based on baseline measures of indicators.
 - If baseline data are not available, use your judgment
- Based on what could be achieved with funding and resources
- Reflect programme strategies tailored to local epidemic
- For two years based on change from baseline
- Often from bio-behavioural surveys.
- Set for impact, outcome, and coverage at the national and sub-national areas



Data Sources for Targets

Use survey data when:	Use programme data when:
Estimating trends or conducting surveillance among the entire population of people who sell sex	Estimating trends among those who receive services
Estimating the proportion of the population reached with services	Estimating the number of people reached with services
Describing the characteristics and behaviours of the population	Describing the characteristics of those reached
Assessing the effectiveness of the STI prevention programme every 2 years	For monthly, quarterly and annual reporting of activities completed
Characterizing those not reached	Characterizing those routinely reached and those newly reached
Assessing whether national and sub-national outcome and coverage targets are reached	Assessing whether service delivery targets are reached

Method	Description
International Reference Method.	Determine whether the baseline estimates are "high" "medium" or "low" based on international guidelines on target setting. (See UNODC Technical Guide for Countries to set targets for universal access to HIV prevention, treatment and care for people who inject drugs). Set the target at the next level higher than the baseline. For example, if the baseline is "low", set a target of "medium."
10 Years to 80% Method.	Identify the baseline indicator: example: 20% and determine the gap between 20% and a target of 80%. If it takes 10 years to get to 80% from 20%, how far can you get in 2 years? The programme should aim to improve 6% each year or 12% in 2 years. At this rate, the target of 80% will be achieved in 10 years. The justification for this approach is that a target of 100% is rarely reached. Targets of 80% are more feasible. Change does not occur quickly. A ten year plan is reasonable for hard to change behaviours. At this rate, the target of 80% will be achieved in 10 years. The justification for this approach is that a target of 100% is rarely reached. Targets of 80% are more feasible. Change does not occur quickly. A ten year plan is reasonable for hard to change behaviours.
X% increase method.	For each indicator, increase the target 10% from baseline. For example, if targets should increase proportionally by 20% and the baseline is 40%, then the target is (20% of 40% = 8%) 48%.

Absolute increase method.	For each indicator, an absolute increase in the baseline of X amount is set as a target. For example, if targets should increase by an absolute 20% and the baseline is 40%, then the target is 60% (40% + 20%= 60%). This type of target setting is often difficult to rationalize.
Expert opinion or consensus.	Some behaviours are harder to change than others and take a longer time to modify. Programmes that are newly implemented may require a longer time to gain the cooperation of the community and see results. Many factors can affect the achievement of targets. In this method, local people including members of the target population assess these factors and set reasonable targets based on their insight and knowledge.
Trends method.	For countries and sub-national areas that have a strong programme and can review trends in indicators, one method is to extend the trend line of each indicator (unless the trend is going the wrong direction!).
Better than the rest method.	If baseline targets are known for several priority prevention areas, targets can be set higher than any area has achieved. The goal is to improve beyond what the best area has achieved. This method is not appropriate in some settings, but may serve to encourage a healthy competition.



Example of Target Setting

X% INCREASE METHOD

For each indicator, increase the target 10% from baseline.

If target is to increase proportionally by 20% annually.

Baseline = 40%, >> target is (20% of 40% = 8%) therefore 40% + 8% = 48%.



Definitions are Key

Defining "reached" can be problematic.

For example:

- •How many condoms does a person need to be "reached" by a condom distribution program? Enough for a day? A week? A month?
- •If one person is picking up condoms for five others, how many have been reached?
- •If a person refuses to get tested for HIV, has he been reached by the testing service? What if he refuses because he knows that he is infected?
- Does seeing a STI prevention poster on a wall meet the definition of being reached with targeted IEC



Service	Simple Definition of "Reached"	More Complete definition of "Reached"
What is the operational definition of "a person reached with a service"? What is the operational definition of a critical enabler intervention being implemented?	Adopt the national definition of "person reached" for service delivery and for reporting	Definition of person reached with a service/ service package



Size Matters (I)

What is the size of the population at risk for STIs in the catchment area?

Why do I care?

To measure coverage – are we doing our jobs?




Size Matters (II)

STEPS:

- Ask 50-100 local people where sex workers are reached.
- Visit and map all places named.
- Sex worker informants will be best




Size Matters (III)






Low Tech Version

Size Matters (IV)





Size Matters (V)



Crude size estimation for service providers

- Count number of SW reached by a service provider in a month. Divide this number by the estimated percent of SW reached by provider.
 - Example: service provider reaches 2000 people/mo.
 Estimates it reaches 50% of all sex workers. There are 4000 sex workers (2000 ÷ .50).
- Estimate the number of people who sell sex at each venue identified during mapping and sum all sites.



Size Matters: Lessons

Request the following information:

- National and sub-national HIV prevalence estimates
- STI prevalence for men and women; by age >< 25
- All reports on STI among SW
- Information on effective HIV/STI prevention programmes
- Information on national HIV/STI prevention targets for most at risk populations
- Important information gaps



Size Matters: Lessons

Sub-groups could include:

- Young people experimenting with drug injection for first time
- Women who sell sex to get money for drugs
- Homeless men who are injecting long-term
- People recently released from prison
- Recent migrants
- Men who have sex with men



Service with a Smile :: Quality Counts

Table II.3.1 Examples of targets for quality, output and coverage indicators

NOTE: Not all possible indicators are included here. Numbers and percentages in the table are examples for illustrative purposes only.

		2010 (baseline)	2011		2012	
Indicator type	Indicator	Baseline estimate, % or N	% change from baseline	Target % or N	% change from baseline	Target % or N
Quality	5.3 Percentage of providers testing and treating for STIs who have been trained to provide STI services to sex workers	60%	+10%	66%	+20%	72%
Output	4.2 Number of sex workers provided with condoms by HIV prevention programmes for sex workers	60	+20%	72	+40%	84
Coverage	3.6 Percentage of sex workers reached by condom promotion and distribution programmes the past 12 months	50%	+20%	60%	+40%	70%



Outputs Monitor

Information needed for specifying output indicators and for monitoring outputs		Examples
1	A well-defined item or activity that can be counted	Number of condoms distributed to clients
2	Whether the quality of the output must be assessed prior to counting it as an output	Whether expired condoms are excluded from the number of condoms distributed
3	A data collection form to obtain information needed to counting the item or activity	An encounter form recording whether condoms were provided to clients of the needle/syringe exchange site
4	A protocol for aggregating the indicator from outreach workers to service center	Computer spreadsheet programme; Hand tally



Outputs Monitor (II)

5	The time period during which activity occurs	Condoms distributed each calendar month
6	Justification that increases in output increases desired outcomes & reduced STI/HIV	Condoms distributed to IDU, if used, reduces probability of sexual transmission among IDU and their sexual partners
7	Baseline estimate of the indicator and feasible target	Baseline: 5000 condoms distributed in January based on providing 10 condoms per month to 500 clients Target: 6000 condoms distributed per month in 2 years. Based on 10 condoms per month to 600 clients
8	Operational definition of what is meaning of "reached"	Reached by a targeted condom distribution programmes is defined as receiving at least two condoms in past calendar month
9	Graphs for displaying trends in the output indicator at the service delivery level	Graph at the service delivery level showing condoms distributed for each calendar month and the target for each month



Double Counting

Method to Avoid Double Counting

Brief Description of Method

Simple

Use when information is not routinely recorded when contacting people who inject drugs

Quarterly Brief Intercept Surveys of Clients during 3-days period

- Ask all contacts accessing the service over 3 days: Is this the first time you have received this service here? If not, when was the last time you received the service?
- Answer: Days ago, weeks ago, months ago, or years ago.

Moderate

Use when information is recorded for each client encounter, but it is not possible to use a Unique Identifier Code

Record whether the encounter is a first time encounter or not

 Ask clients if the contact with the service is the first one this month, first one this year, or first ever

Complete

Use when it is feasible to collect a Unique Identifier Code and to link records

For each encounter, record the service provided and the individual's Unique Identifier Code Record each service provided to the client at the time the service is provided using a Unique Identifying Code (UIC) to link services provided to the same client over time



Double Counting Solved: Simple

Quarterly Brief Intercept Surveys of Clients during 3-days period

- Ask all contacts accessing the service over 3
 days: Is this the first time you have received
 this service here? If not, when was the last
 time you received the service?
- Answer: Days ago, weeks ago, months ago, or years ago.



Double Counting Solved: Moderate

Record whether the encounter is a first time encounter or not

 Ask clients if the contact with the service is the first one this month, first one this year, or first ever



Double Counting Solved: Complete

For each encounter, record the service provided and the individual's Unique Identifier Code Record each service provided to the client at the time the service is provided using a Unique Identifying Code (UIC) to link services provided to the same client over time



Quality Assessments: A Checklist

STI clinics

- □ Clients receive IEC when accessing STI clinc
- Clients receive condoms when accessing STI clinic
- □ Clients' risk assessed for comprehensive testing

Outreach interventions

- Clients receive sufficient condoms to protect all sex acts
- ☐ Clients receive STI/HIV testing referrals
- □ Clients are engaged voluntarily



Quality Assessments: A Checklist

Quality Checklist for all Services:

There is a system in place to ensure no stock-outs occur
UIC or other system exists to count the number of unique clients rather than the number of contacts with the service
There is an established referral system including a follow-up mechanism
IEC is provided
A risk assessment is conducted
Condoms are provided for sexually active IDU



Data collection and Management

 Must I use a unique identifier? How do I make a data collection form? Pencil and paper vs. digital?



Data Management

Encounter Form

(see Tools x, y)

- Records the Unique Idenitfier Code, and the characteristics of the person reached by the service delivery organisation
- Records the services provided
- Can be simple or comprehensive

Referral Forms

(see Tools x, y)

- Records that a referral for a service was made
- Records the result of the referral

Aggregation Forms

(see Tools x, y)

 Used by service delivery providers to aggregate data from individual encounter forms



Sample Encounter Form

A1 Type of Location	A3 Today's Date Day/ Month/ Year
 A1 Type of Location 1 STI clinic site (fixed) 2 Condom distribution site (outreach) 3 Other provider (specify	A4 Service delivery provider (eg. NGO) number
A2 Sub-national area number	A5 ID number of individual providing service:

B1 Unique Identifier Code	B5 Gender: 1 M 2 F 3TG-MTF 4TG-FTM	
B2 First EVER visit to site? 1 YES 2 NO	B6 Birth Date Day/ Month/ Year	
B3 First visit since January 1: 1 YES 2 NO	B7 Primary Language:	
B4 Other Service Providers Visited in Past 30 Days:	B8 When Moved to This Area: Month:Year	



Drugs Used in past 30 Days	A Smoked/ Sniffed/ Other	B Inject		Last	C time, Clean Needle ?	D Times Injecte d Past 7 Days	E Of those, times with Clean Needle	F Client Stage for Using Clean Needles in Past 7 Days
C1 Cocaine	1	Υ	N	Υ	N			
C2 Crack	1	Y	N	Υ	N			
C3 Heroine	1	Y	N	Υ	N			1 Did not consider 2 Thought about it 3 Tried First Time 4 Inconsistent Use 5 Consistent Use
C4 Speed/Meth	1	Y	N	Y	N			
C5 Other Opioids	1	Y	N	Y	N			
C6 Other (1)	1	Y	N	Y	N			
C7 Other (2)	1	Y	N	Υ	N			
C8 Sex in Past 30 Days	A Sex for Mon ey		er of Partn		me, Jsed Condo n?	D Times Sex In Past 7 Days	E Of those, times with Condom	F Client Stage for Condom Use in Past 7 Days (Use above codes)



Referral Card and Form for Monitoring Referrals

Date Referred:			
UIC	Referred by (A4)	Referred to (Code)	Service Requested (E)



Referral Follow Up

Date Referre d	UIC	Service Requested (code from E):
Referral Results (check one):	 Person never came to referral site Person came to referral site – doesn't need to return Person came to referral site needs to return Person came –referred elsewhere 	Date seen: Date referred:



Data use/reporting

I have piles of data. What do I do with it?



Data use/reporting

- Don't be overwhelmed
- 2. Ask simple, answerable, and actionable questions E.g., Did I have a condom stock out in the past 6 months?
- 3. Ask for help
- Put analysis and reporting in budget and work plan



Reporting Loops

