

Updates on Contraceptive Technology Part 2

Mario Festin

Human Reproduction Team

Department of Reproductive Health and Research

Twitter [@HRPresearch](#)



Key Facts about family planning/contraception

- ❑ 214 million women of reproductive age in developing countries who want to avoid pregnancy are not using a modern contraceptive method (unmet need for modern contraception).
- ❑ Some family planning methods, such as condoms, help prevent the transmission of HIV and other sexually transmitted infections.
- ❑ Family planning / contraception reduces the need for abortion, especially unsafe abortion.
- ❑ Family planning reinforces people's rights to determine the number and spacing of their children.
- ❑ By preventing unintended pregnancy, family planning /contraception prevents deaths of mothers and children.

Main benefits of family planning/contraception

- ❑ Secures the well being and autonomy of women
- ❑ Supports the health and development of communities
- ❑ Prevents unplanned pregnancy and pregnancy-related health risks of women
- ❑ Prevents adolescent pregnancy
- ❑ Reduces infant mortality and prevents HIV/AIDS transmission to newborns
- ❑ Empowers people and enhances education
- ❑ Slows population growth

Unmet need for contraception

Definition

- Women with unmet need are those who are **fecund and sexually active** but are **not using any method of contraception**, and report **not wanting any more children or wanting to delay the next child**.
- The concept of unmet need points to the gap between women's **reproductive intentions** and their **contraceptive behavior**.

Unmet need is especially high among groups such as.


- Adolescents
- Migrants
- Urban slum dwellers
- Refugees
- Women in the postpartum period

Reasons for unmet need for modern FP/ C

- ❑ limited choice of methods;
- ❑ limited access to contraception, particularly among young people, poorer segments of populations, or unmarried people;
- ❑ fear or experience of side-effects;
- ❑ cultural or religious opposition;
- ❑ poor quality of available services;
- ❑ users and providers bias;
- ❑ gender-based barriers.

World contraceptive use





Percentage of women using some method of contraception among those aged 15-49 who are married or in a union



Comparing Effectiveness of Family Planning Methods

More effective

Less than 1 pregnancy per 100 women in one year



Less effective

About 30 pregnancies per 100 women in one year

How to make your method more effective

Implants, IUD, female sterilization:

After procedure, little or nothing to do or remember

Vasectomy: Use another method for first 3 months

Injectables: Get repeat injections on time

Lactational Amenorrhea Method (for 6 months):

Breastfeed often, day and night

Pills: Take a pill at the same time each day

Patch, ring: Keep in place, change on time

Condoms, diaphragm: Use correctly every time you have sex

Fertility awareness methods: Abstain or use condoms on fertile days. Standard Days Method and Two-Day Method may be easier to use.

Withdrawal, spermicides: Use correctly every time you have sex

Outline and objectives

- ❑ Description of the method
- ❑ Mechanism of action
- ❑ Effectiveness
- ❑ Eligibility criteria
- ❑ Benefits and side effects
- ❑ Interventions for associated effects

Methods

- ❑ Combined oral contraceptives
- ❑ Injectable contraceptives, progestin-only
- ❑ Injectable contraceptives, combined
- ❑ Hormonal implants


- ❑ **IUDs (copper bearing)**
- ❑ **LNG IUS**
- ❑ **Male and female condoms**
- ❑ **Other barrier methods**
- ❑ **Fertility awareness, lactational amenorrhea**
- ❑ **Emergency contraception**
- ❑ **Tubal ligation and vasectomy**
- ❑ **Other methods**

Copper Intrauterine Contraceptive Devices (IUDs or IUCDs)




Copper T 380A

What is a Copper IUD?



- Small plastic device with copper sleeves or wire around it, inserted through the vagina and cervix into the uterus
 - Safe
 - Highly effective
 - Long acting (up to 12 years)
 - Require trained provider to insert and remove

Copper-bearing IUD



Copper T-380A

TCu-380A, “Copper T” is most widely used copper IUD




Multiload 375

Multiload 375 – less commonly available

Copper T: Mechanism of Action

Prevents fertilization by:

- ❑ Impairing the viability of the sperm
- ❑ Interfering with movement of the sperm



Source: Ortiz, 1996

Copper IUDs: Characteristics

- ❑ Safe and highly effective
- ❑ Require no user action
- ❑ Long-acting (up to 12 years)
- ❑ Rapid return to fertility
- ❑ No systemic effects
- ❑ Other health benefits
- Potential side effects
- Require pelvic exam
- Trained provider needed to insert and remove
- Possible pain or discomfort during insertion
- Do not protect against STIs/HIV
- Rare complications

Source: CCP and WHO, 2011

Copper IUDs: Possible Side Effects

Some women may experience:

- ❑ More cramps and pain during menstruation
- ❑ Increased or prolonged menstrual bleeding
- ❑ Bleeding between menstrual periods



Side effects are most common during the first 3–6 months.

Source: CCP and WHO, 2011; Larsson, 1993; DeMaeyer, 1989; WHO, 2004, updated 2008; WHO Special Programme of Research Development and Research Training in Human Reproduction, 1997.

Copper IUDs: Health Benefits

IUDs are known to:

- ❑ Reduce risk of ectopic pregnancy
 - Rate in all IUD users is 12 in 10,000 (2 in 10,000 for Copper T380A users)
 - Rate in women using no contraception is 65 in 10,000
- ❑ Help protect against endometrial cancer

Source: CCP and WHO, 2011; Sivin, 1991.

Potential Complications

- ❑ Perforations—Very rare, 1 in 1,000 insertions
 - Linked to skill and experience of provider
 - Reduced through supervised training
- ❑ PID—Rare, most due to gonorrhea or chlamydia at time of insertion
- ❑ Expulsions—Related to provider skill, age and parity of woman and insertion factors

Expulsion Rates Are Higher for Postpartum Insertion

Timing of Insertion	Expulsion Rates
Interval (more than 6 weeks after delivery)	Low (3% for skilled provider)
Immediate postpartum (within 10 minutes)	Slightly higher
Early postpartum (between 10 minutes and 48 hours)	Moderately higher
Late Postpartum (48 hours to 4 weeks)	High - Generally not recommended

Source: Chi, et al, 1985.

Who Can Initiate Copper IUDs

WHO MEC category 1 and 2 examples

WHO MEC Category	Conditions (selected examples)
Category 1	≥20 years, cervical ectopy, uterine fibroids without distortion of the uterine cavity, irregular bleeding without heavy bleeding, breastfeeding > 6 weeks, history of PID
Category 2	Menarche to <20 years, nulliparous, heavy or prolonged bleeding, severe dysmenorrhea, anemia, HIV/AIDS infection, with or without ARV therapy

Source: WHO, 2004; updated 2008.

Who Should Not Initiate Copper IUDs

WHO MEC category 3 and 4 examples

WHO MEC Category	Conditions (selected examples)
Category 3	48 hours to <4 weeks postpartum, ovarian cancer/if initiating use, high individual risk of STI/ if initiating use
Category 4	pregnancy, unexplained vaginal bleeding (prior to evaluation), current PID or cervical infection, endometrial or cervical cancer/if initiating use

Source: WHO, 2004; updated 2008.

Timing of IUD Insertion

Interval insertion

- ❑ Within the first 12 days of menstrual cycle
- ❑ Any other time if woman is not pregnant

Postpartum insertion

- ❑ Immediately after vaginal or cesarean delivery if no infection or bleeding complications
- ❑ Within 48 hours or delay at least 4 weeks

Postabortion insertion

- ❑ Immediately or within 12 days if no infection

Source: WHO, 2004; updated 2008.

Side Effects

- ❑ Common side effects:
 - Heavier and/or prolonged menstrual bleeding
 - Menstrual cramping
 - Spotting between periods

- ❑ Side effects:
 - Are not signs of illness
 - Usually become less within the first 3–6 months

Correcting Misconceptions

IUDs:

- Rarely lead to PID
- Do not increase risk of STIs, including HIV
- Do not work by causing abortion
- Do not make women infertile
- Do not move to the heart or brain
- Do not cause birth defects
- Do not cause pain for either partner during sex
- Significantly reduce risk of ectopic pregnancy

Source: CCP and WHO, 2011.

Signs of Possible IUD Complications

- Bleeding and severe abdominal cramping within a few days post-insertion → perforation
- Irregular bleeding or pain every cycle → partial expulsion, perforation
- Fever, unusual vaginal discharge, low abdominal pain → infection
- Missing IUD strings, missed period → expulsion, pregnancy

Source: CCP and WHO, 2011.

Heavy, Prolonged or Irregular Bleeding

Counseling and reassurance are key

Problem	Action/Management
Heavy or prolonged bleeding	<ul style="list-style-type: none">● Reassure client that this is common and not harmful, usually diminishes after few months● For short-term relief offer 5-day course of tranexamic acid or NSAIDs (not aspirin)● Provide iron tablets
Irregular bleeding	<ul style="list-style-type: none">● Reassure client● For short-term relief offer ibuprofen or indomethacin 2 times daily after meals for 5 days

Managing IUD Side Effects or Complications: Cramping and Mild Pain

Counseling and reassurance are key

Problem	Action/Management
Cramping and mild pain	<ul style="list-style-type: none">● She can expect cramping and pain in first 1–2 days after insertion● Reassure client that this is common in first 3–6 months, not harmful, usually decreases over time● Suggest ibuprofen, other pain reliever (not aspirin if she also has heavy bleeding)● If cramping continues, occurs outside of menstruation, evaluate, treat or refer● If cramping is severe but no underlying condition, discuss removing the IUD

Source: CCP and WHO, 2011.

Managing IUD Side Effects or Complications:

Severe Pain in Lower Abdomen (Rare)

- ❑ Rule out PID, ectopic pregnancy or perforation.
- ❑ If PID is suspected, treat with appropriate antibiotics for gonorrhoea, chlamydia and anaerobic bacterial infection. There is no need to remove the IUD.
- ❑ If ectopic pregnancy is suspected, refer immediately.

Source: CCP and WHO, 2011.

Managing IUD Side Effects or Complications: Suspected Perforation

- ❑ Stop procedure immediately, remove IUD
- ❑ Observe vital signs for an hour; check for signs of bleeding
 - If rapid pulse, falling blood pressure, or increased pain: refer
- ❑ Provide alternative contraception
 - Advise avoid sex for 2 weeks
- ❑ Follow-up in a week or as needed

Source: CCP and WHO, 2011.

Missing Strings

- ❑ Determine risk of pregnancy
- ❑ Perform pelvic exam, probe for strings in cervical canal
- ❑ If cannot locate strings, consider X-ray or ultrasound, or refer
- ❑ Give choice of another contraceptive method
- ❑ Insert another IUD if expulsion is confirmed and
 - Woman is not pregnant
 - She still wants to use an IUD

Source: CCP and WHO, 2011.

Suspected Pregnancy

- ❑ Assess for pregnancy, including ectopic pregnancy
- ❑ If the client is pregnant and wishes to continue the pregnancy:
 - Explain that using an IUD during pregnancy increases the risk of preterm delivery or miscarriage
 - If possible, remove the IUD
 - If not possible to remove, advise close follow-up for signs of septic miscarriage

Hormonal intrauterine device



LNG-IUS

Hormonal IUDs



- ❑ Steadily release a progestin
- ❑ Levonorgestrel IUD (LNG-IUD) most common hormonal IUD
- ❑ Also known as LNG-intrauterine system (LNG-IUS)
- ❑ Immediately reversible
- ❑ Effective for 5 years
- ❑ Category 4 if immediate post-septic abortion, before evaluation of unexplained vaginal bleeding, or with malignant trophoblastic disease, with current breast cancer

Other features of the LNG IUS

- ❑ Works primarily by suppressing the growth of the lining of the uterus
- ❑ Needs to be inserted into a uterus by a trained health care provider


- ❑ Some report lighter, fewer or even absent bleeding days
- ❑ Also infrequent reports of headaches, breast tenderness or pain, acne

Male and Female Condoms




What Is a Male Condom?

- ❑ A sheath that fits over a man's erect penis
- ❑ Most are made of thin latex rubber
- ❑ Other materials include natural skin and various synthetics



What Is a Female Condom?

- A transparent sheath that fits loosely inside a woman's vagina
- Most common type is FC2 —made of thin, soft, synthetic rubber film
 - Has flexible rings at both ends
- Also natural latex female condoms (Cupid) available in some countries




Comparing Effectiveness of Male and Female Condoms

In real-life situations, correct and consistent use may be difficult to achieve

Pregnancy rates:

Male



Female




perfect use	2%	5%
typical use	15%	21%

Source: Hatcher, 2007.


Condoms: Mechanism of Action

- ❑ **Create a barrier that keeps sperm from entering the vagina, thus preventing pregnancy**
- ❑ **Also form a barrier against STIs including HIV**



Male and Female Condoms

Dual Protection




Condoms are the *only* FP method that provides **dual protection**:

- ❑ Protection from pregnancy *and*
- ❑ Protection from transmission of HIV and other STIs between partners

Effectiveness for Preventing HIV and STIs

- ❑ The consistent, correct use of condoms significantly reduces the risk of HIV infection in men and women
- ❑ When used correctly with every act of sex, condoms are 80%–95% effective in preventing HIV infection
- ❑ Condoms prevent STIs transmitted through body fluids such as gonorrhea and chlamydia
- ❑ Condoms are less effective for preventing STIs transmitted by skin-to-skin contact, such as herpes and warts

Characteristics of Male Condoms



- ❑ Safe and easy to use
- ❑ Widely available
- ❑ Effective when used consistently and correctly
- ❑ Provide dual protection
- ❑ No hormonal side effects
- ❑ Can help men with premature ejaculation
- ❑ Do not require provider's help
- ❑ Can be used as temporary backup method
- ❑ Protect women from conditions caused by STIs


Characteristics of Male Condoms

- ❑ As typically used, less effective than many other methods
- ❑ Require partner communication and cooperation
- ❑ Can be damaged by oil-based lubricants, heat, humidity or light
- ❑ May reduce sensation
- ❑ Can interrupt sex





Who Can Use Male Condoms

- All men and women can safely use male condoms, *except* those with severe allergic reaction to latex
 - Extremely rare among both men and women
 - Non-latex condoms are available in some countries



Managing Problems

- If condom not used—Offer emergency contraception
- If slipping or breaking—Ask about practices, behaviors
- If difficulty putting on—Ask client to demonstrate, correct errors
- Difficulty persuading partner to use—Help her choose approaches that will work
Suggest adding another method



Managing Problems

Mild irritation or mild allergic reaction to condom—Itching, redness, rash and/or swelling

- Try another brand of condoms
- Put lubricant or water on the condom
- If symptoms continue, assess or refer for possible vaginal infection or STI
- If no infection, may have allergy to latex

What Men and Women Like

What Women Like:

- ❑ Female-controlled
- ❑ Texture feels more natural than latex male condoms
- ❑ Offer STI/HIV protection
- ❑ Outer ring provides stimulation
- ❑ Do not require provider's help

What Men Like:

- ❑ Can be inserted in advance
- ❑ Are not tight or constricting
- ❑ Do not dull sensation
- ❑ Do not have to be removed immediately

What Men and Women Don't Like

Limitations:


- ❑ May be difficult to insert at first, require practice
- ❑ Not as effective as other methods
- ❑ More expensive than male condoms
- ❑ Less available than male condoms

Who Can Use Female Condoms

- All men and women can use FC2
- Latex female condoms: Severe allergic reaction to latex is the only condition that prevents use
 - Extremely rare

Managing Problems

- If having trouble inserting: Ask her to demonstrate
- If uncomfortable: Reinsert or reposition
- If condom squeaks: Use more lubricant
- If condoms slips or is not used correctly: Offer emergency contraception
- Difficulty persuading partner: Help her choose approaches that will work



Managing Problems

(Continued)

- ❑ Mild irritation in or around the vagina or penis (itching, redness, rash)
 - Usually goes away on its own
 - Suggest added lubricant inside condom or on penis
 - If symptoms persist, assess and treat for possible vaginal infection or STI

Responding to Myths and Fears

Male condoms:

- ❑ Do not make men impotent, weak, or decrease sex drive
- ❑ Do not cause illness in men because sperm “back-up”

Female condoms:

- ❑ Are not difficult to use, but correct use needs to be learned
- ❑ Do not cause illness in women because they prevent semen from entering the body

Responding to Myths and Fears

(continued)

Both male and female condoms:

- ❑ Cannot get lost in woman's body
- ❑ Do not have holes that HIV can pass through
- ❑ Are not intended only for use outside of marriage, but also used by married couples

Other barrier methods

- ❑ **Diaphragm** – is placed deep in the vagina before sex, covering the cervix as a barrier, usually used with a spermicide
- ❑ When used correctly with every act of sex, 6 pregnancies per 100 women using the diaphragm over the first year.
- ❑ **Cervical caps** – is also placed deep in the vagina before sex, covering the cervix as a barrier.
- ❑ Not as effective, with 20 pregnancies per 100 women in the first year.



Other barrier methods

- ❑ Spermicide – applied deep in the vagina before every act of sex
 - Usually Nonoxynol 9, but other substances also available
 - Work by causing the membrane of sperm cells to break, killing them or slowing movement.
 - One of least effective methods, may be a primary or a back up method.
- ❑ Microbicide (either anti-infective alone or dual protection)



Fertility awareness methods


- ❑ Requires the cooperation of both partners, with a commitment to abstain or use another method on fertile days
 - Periodic abstinence
 - Natural family planning
- ❑ Must be aware of the body changes or keep track of days, according to rules of the specific methods.
- ❑ No side effects.

Fertility awareness methods


- ❑ Calendar based methods
 - Standard days methods
 - Calendar rhythm methods
- ❑ Symptoms based methods
 - Cervical secretions
 - Basal Body Temperature
 - Increase noted slightly after release of the egg or ovulation
 - Two day method
 - Sympto-thermal method

Standard Days Methods

- ❑ 95% effective with consistent and correct use
- ❑ A woman can use the SDM if most of her menstrual periods are 26 to 32 days long.
- ❑ Days 8 to 19 of each cycle are considered as fertile days.
- ❑ The couple can use cycle beads, a color-coded string of beads that indicates fertile and nonfertile days.



How to use cycle beads



- Move ring to RED bead when period starts

1



- Move ring to next bead every day. Move ring even on bleeding days

2



- Use condoms or abstain when ring is on WHITE beads

3




- BROWN beads are safe days of no pregnancy

4



- When period starts again move ring to red bead to begin again.

5



- Always check your period comes between dark brown bead and last brown bead.



Lactational amenorrhea

- ❑ A family planning method based on breastfeeding
- ❑ Can be effective when all the following conditions are present
 - Infant is less than 6 months old,
 - as long as monthly bleeding has not returned and
 - the woman is fully breastfeeding.
- ❑ Provides an opportunity to offer a woman an ongoing method for continuously 6 months

Lactational amenorrhea

- ❑ A woman with the following may want to consider other methods:
 - Has HIV infection
 - Especially if the woman is not taking ARVs, not fully breastfeeding, newly infected or has advanced disease.
 - Using certain medications (mood-altering drugs, reserpine, ergotamine, anti-metabolites, cyclosporine, high dose corticosteroids, bromocriptine, radioactive drugs, lithium and certain anticoagulants)
 - A newborn with a condition that makes it difficult to breastfeed

Emergency contraception

- ❑ Emergency contraceptive pills help to prevent pregnancy, if taken up to 5 days after unprotected sex and other situations such as
 - forced sex (rape) and contraceptive mistakes (condom slippage or breakage, misplaced IUD, late for injections, etc.)
- ❑ Do not disrupt an existing pregnancy
- ❑ Safe for all women
- ❑ Provide an opportunity for women to start using an ongoing FP method
- ❑ Effectiveness (no method = 8 pregnancies/100 woman years)
 - Progestin only ECPs – 1 pregnancy per 100 women years
 - Combined estrogen progestin ECPs – 2 pregnancies per 100 women years.

Emergency contraception

- ❑ Contains a progestin alone (levonorgestrel or ulipristal) or a progestin and an estrogen together
 - ❑ Works primarily by preventing or delaying ovulation
 - ❑ Does not work if the woman is already pregnant.
-
- ❑ Copper IUD effective as emergency contraception, when inserted up to 7 days after unprotected sex
 - ❑ IUDs when used as emergency contraception, can then be continued to be regular contraception

Pill Type and Hormone	Formulation	Pills to Take		
		At First	12 Hours Later	
Dedicated ECP Products				
Progestin-only	1.5 mg LNG	1	0	
	0.75 mg LNG	2	0	
Ulipristal acetate	30 mg ulipristal acetate	1	0	
Oral Contraceptive Pills Used for Emergency Contraception				
Combined (estrogen-progestin) oral contraceptives	0.02 mg EE + 0.1 mg LNG	5	5	
	0.03 mg EE + 0.15 mg LNG	4	4	
	0.03 mg EE + 0.15 mg LNG	4	4	
	0.03 mg EE + 0.125 mg LNG	4	4	
	0.05 mg EE + 0.25 mg LNG	2	2	
	0.03 mg EE + 0.3 mg norgestrel	4	4	
	0.05 mg EE + 0.5 mg norgestrel	2	2	
	Progestin-only pills	0.03 mg LNG	50*	0
		0.0375 mg LNG	40*	0
	0.075 mg norgestrel	40*	0	

* Many pills, but safe. See Question 8, p. 63.
LNG = levonorgestrel EE = ethinyl estradiol

Family Planning Global Handbook

Correcting misconceptions about Emergency contraception


ECPs :

- ❑ Do not cause abortion
- ❑ Do not cause birth defects if pregnancy occurs
- ❑ Are not dangerous to a woman's health
- ❑ Do not promote sexual risk taking
- ❑ Do not make women infertile

Female Sterilization


Tubal Ligation

- ❑ Permanent method, done through mini-laparotomy (small incision) or laparoscopy
- ❑ One of the most effective methods with less than 5 failures per 1000 women
- ❑ May be used by any woman
- ❑ Counselling needed to ensure no post procedural regrets



Male Sterilization

Vasectomy



- ❑ Simple permanent surgical method in men
- ❑ With a 3 month delay in taking effect due to storage of sperm
- ❑ Tubes that carry sperm to the penis (vas deferens) are cut and ligated (by sutures or cautery)
- ❑ May be done using non-scalpel technique
- ❑ Less than 2 per 1000 pregnancies over first year
- ❑ Does not affect male performance nor provides increase risk for cancer

Other methods

□ Ring

- Requires keeping a flexible ring which releases progestins and estrogen which are absorbed by the body
- Kept for 3 weeks, and woman menstruates in 4th week
- Prevents ovulation
- Progesterone only vaginal rings may be used by breast feeding women

□ Patch

- Requires wearing a small adhesive patch with estrogen and progestins weekly, for 3 weeks
- Works by preventing ovulation




Acknowledgments and References

- ❑ Main Reference:
 - Family Planning – A Global Handbook for Providers (<https://www.fphandbook.org/>)
- ❑ Acknowledgements
 - Family Health International
 - Knowledge for Health
 - Institute of Reproductive Health

MEC 5th Edition Recommendations

- Details of the recommendations on the methods are available in the full MEC 5th edition document.
- http://www.who.int/reproductivehealth/publications/family_planning/MEC-5/en/



Follow us on Twitter [@HRPresearch](https://twitter.com/HRPresearch)

Visit our website who.int/reproductivehealth

Email us at: hrx-info@who.int