The WHO postpartum family planning compendium

Petrus Steyn Scientist, WHO/ SRH/ CFC

Twitter **@HRPresearch**





WHO Guidance on Postpartum Family Planning





Guide women through their postpartum family planning options

The Postpartum Family Planning Compendium integrates core WHO guidance for clinicians, program managers, and policy-makers.







The WHO Postpartum Family Planning Compendium


CASE STUDY



Yesterday, a 32-year-old woman delivered a healthy baby 38 weeks gestation at the local hospital. She is **breastfeeding**. She has a history of **hypertension and smokes**. She is interested in the contraceptive implant. *Is she eligible for the contraceptive implant prior to her discharge from the hospital?*

Choose Medical Conditions:	Medical Conditions: Breastfeeding Status:		Current Medication Use:		Other	Other Characteristics:	
0 Selected	O Yes	O No	0 Selecte	t	0 Sel	ected	
• Time interval after delivery:		hours	3 weeks	4 weeks	6 weeks	6 months	12 months
	-						-

Yesterday, a 32-year-old woman delivered a healthy baby 38 weeks gestation at the local hospital. She is breastfeeding. She has a history of hypertension. She is interested in the contraceptive implant. Is she eligible for the contraceptive implant prior to her discharge from the hospital?



Yesterday, a 32-year-old woman delivered a healthy baby 38 weeks gestation at the local hospital. She is breastfeeding. She has a history of hypertension. She is interested in the contraceptive implant. Is she eligible for the contraceptive implant prior to her discharge from the hospital?




Yesterday, a 32-year-old woman delivered a healthy baby 38 weeks gestation at the local hospital. She is breastfeeding. She has a history of hypertension. She is interested in the contraceptive implant. Is she eligible for the contraceptive implant prior to her discharge from the hospital?




Yesterday, a 32-year-old woman delivered a healthy baby 38 weeks gestation at the local hospital. She is **breastfeeding**. She has a history of **hypertension**. She is interested in the contraceptive implant.

Is she eligible for the contraceptive implant prior to her discharge from the hospital?









Contraceptive implant, within 48 hours postpartum



Description

Small, flexible rods or capsules placed under the skin of the upper arm; contains progestogen hormone only

Effectiveness to prevent pregnancy

How it works

Thickens cervical mucous to block sperm and egg from meeting and prevents ovulation

Comments

Implants: Health-care provider must insert and remove; can be used for 3–5 years depending on implant, irregular vaginal bleeding common but not harmful



Contraceptive implant, within 48 hours postpartum

Conditions / Subconditions

Breastfeeding Status:

Breastfeeding	Category	Clarifications/Special consideration
	LNG /ETG	
a) < 6 weeks postpartum	2	
b) ≥ 6 weeks to < 6 months postpartum (primarily	Ť.	
breastfeeding)	1	
c) ≥ 6 months postpartum		


Medical Condition:

Hypertension	Category	Clarifications/Special consideration
	LNG /ETG	
a) History of hypertension, where blood pressure CANNOT be evaluated (including hypertension in pregnancy)	2	Read More
b) Adequately controlled	1	



Guide women through their postpartum family planning options

The Postpartum Family Planning Compendium integrates core WHO guidance for clinicians, program managers, and policy-makers.



2. Click on family planning method to see recommendations



Postpartum family planning guidance at your fingertips: Place a bookmark!

https://postpartumfp.srhr.org/





Guide women through their postpartum family planning options

The Postpartum Family Planning Compendium integrates core WHO guidance for clinicians, program managers, and policy-makers.

Choose Medical Conditions:	Breastfeeding Status:	Current Medication Use:	Other Characteristics:
	O Yes O No		



Thank You!

Follow us on Twitter **@HRPresearch**

Visit our website

https://www.who.int/teams/sexual-and-reproductive-health-and-research-(srh)/

